

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/319296919>

Drop Jump: A Technical Model for Scientific Application

Article in *Strength and Conditioning Journal* · August 2017

DOI: 10.1519/SSC.0000000000000331

CITATIONS

42

READS

20,690

5 authors, including:



Jason Pedley

Cardiff Metropolitan University

20 PUBLICATIONS 124 CITATIONS

[SEE PROFILE](#)



Rhodri Lloyd

Cardiff Metropolitan University

158 PUBLICATIONS 5,665 CITATIONS

[SEE PROFILE](#)



Paul Read

University College London

131 PUBLICATIONS 2,857 CITATIONS

[SEE PROFILE](#)



Isabel S Moore

Cardiff Metropolitan University

73 PUBLICATIONS 1,136 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



The influence of muscle architecture on stretch-shortening cycle performance in youth [View project](#)



Injury Prevalence across Male and Female Age-Group Athletes in the international youth Field hockey environment and its ability to influence a best practice 'Pre-habilitation' resource to mitigate Injury risk. [View project](#)

Drop Jump: A Technical Model for Scientific Application

Jason S. Pedley, MSc,¹ Rhodri S. Lloyd, PhD, CSCS*D,^{1,2,3} Paul Read, PhD, CSCS*D,^{4,5} Isabel S. Moore, PhD,⁶ and Jon L. Oliver, PhD^{7,2}

¹Youth Physical Development Centre, Cardiff Metropolitan University, Cardiff, United Kingdom; ²Sports Performance Research Institute New Zealand, AUT University, Auckland, New Zealand; ³Centre for Sport Science and Human Performance, Waikato Institute of Technology, Waikato, New Zealand; ⁴Athlete Health and Performance Research Centre, Aspetar Orthopaedic and Sports Medicine Hospital, Doha, Qatar; ⁵Youth Physical Development Group, School of Sport, Cardiff Metropolitan University, Cardiff, United Kingdom; ⁶Cardiff School of Sport, Cardiff Metropolitan University, Cardiff, United Kingdom; and ⁷Youth Physical Development Group, Cardiff Metropolitan University, Cardiff, United Kingdom

ABSTRACT

PLYOMETRIC TRAINING HAS NUMEROUS BENEFITS TO ATHLETIC DEVELOPMENT. THE DROP JUMP IS A CORNERSTONE EXERCISE FOR DEVELOPING FAST STRETCH-SHORTENING CYCLE FUNCTION. THE DROP JUMP CAN ALSO BE USED AS A READINESS-TO-TRAIN MARKER IN PRACTICAL SETTINGS. TO ASSIST COACHES WITH ATHLETE DEVELOPMENT, A TECHNICAL MODEL FOR THE DROP JUMP IS PROPOSED. VERBAL INSTRUCTIONS PROVIDED BY THE COACH HAVE AN EFFECT ON THE KINETICS AND KINEMATICS OF A TASK PERFORMED BY AN ATHLETE. TO ASSIST COACHES IN FOSTERING EFFECTIVE PERFORMANCE ADAPTATIONS. EXAMPLE EXTERNAL CUES AND ANALOGIES ARE PROVIDED TO HELP WITH OPTIMIZING PERFORMANCE DEVELOPMENT WHILE MINIMIZING INJURY RISK.

INTRODUCTION

Plyometric training is a commonly used method for developing a variety of athletic qualities including speed strength, sprinting speed, explosive power, and running economy (10,26–29,38,79,90). Such training seeks to exploit the force potentiating capabilities of the stretch-shortening cycle (SSC) to improve athletic performance (47). The SSC consists of (a) a rapid eccentric muscle action followed by (b) an isometric amortization phase, and (c) a concentric muscle action (12). This sequence results in an enhanced concentric force output through a number of mechanisms including utilization of elastic energy and reflex muscle activity (47). Most lower-limb plyometric exercises involve jumping, often incorporating the need to rebound against the floor in an attempt to ensure short ground-contact times (<250 ms) (75). The drop jump is a commonly used fast SSC plyometric exercise among strength and conditioning coaches (13,54). In addition to enhancing performance, the drop jump is also used as a readiness-to-train monitoring tool (56) and for injury-risk screening purposes (64,71,76).

There is a large body of research available on plyometric training (6,7,23) and a growing volume of evidence

investigating effective cueing strategies for athletic training (24,53,93,94). However, despite widespread use of the drop jump in strength and conditioning programs, there remains a lack of consensus regarding the correct technical model and effective coaching strategies to improve an athlete's execution of the exercise. A sound technical model will assist in coaching athletes to perform drop jumps in a safe and proficient manner. The likely outcomes of such practice should be a reduction in injury frequency and enhanced physical performance. Two drop jump techniques are defined in peer-reviewed literature: the bounce drop jump and the countermovement drop jump (13). The countermovement drop jump involves a large downward movement during the ground-contact phase. By contrast, the bounce drop jump seeks to immediately reverse downward velocity on landing to minimize ground-contact time. The origins of the first of these 2 derivatives can be traced back to the work of Verkoshansky (86), who describes a “depth jump” exercise where

KEY WORDS:

coaching; injury prevention; plyometric; stretch-shortening cycle

Address correspondence to Jason S. Pedley, jpedley@cardiffmet.ac.uk.

the athlete steps off a box and then performs a vertical jump for maximal height on landing. No rigid restrictions are placed on the magnitude of leg flexion or ground-contact time, although the exercise should be performed quickly. The target outcome of this training method was the development of explosive and maximum strength through stimulation of the central nervous system as a result of the impact. The second variant originates from the work of Komi and Bosco (48) who describe a “drop jump” where the athlete

drops from a platform and immediately executes a vertical jump on ground contact. Emphasis is placed on a short ground-contact time with low magnitudes of leg flexion. The target outcome of this exercise is fast SSC development from the muscle-tendon units of the leg extensors. However, through the passage of time, these 2 exercises have become confused and now many textbooks, authors, and coaches use the terms depth jump and drop jump synonymously (11,21,40,42), to indicate different exercises (22) or to indicate a single

exercise with variations in execution (77). It is proposed that nomenclature should be henceforth standardized with a “drop jump” being an exercise involving restricted amplitude of leg flexion during ground contact and a “depth jump” having unrestricted levels of leg flexion and subsequently less emphasis placed on short ground-contact times as demonstrated in Figure 1. This article will propose a technical model for the drop jump that allows athletes to maximize potential training adaptations. Example cues are provided




Figure 1. Depth jump involves larger amplitudes of center of mass displacement during ground contact than the drop jump. Consequently, the drop jump requires much higher levels of leg stiffness. Depth jumps allow greater jump height to be achieved, but this is at the expense of ground-contact time and therefore, although this exercise might be suitable for promoting some desirable adaptations, it is not suitable for developing fast SSC ability (<250 ms). SSC = stretch-shortening cycle. Depth jump (A) involves larger amplitudes of center of mass displacement during ground contact than the drop jump (B). Consequently, the drop jump requires much higher levels of leg stiffness. Depth jumps allow greater jump height to be achieved, but this is at the expense of ground contact time and therefore, although this exercise might be suitable for promoting some desirable adaptations, it is not suitable for developing fast SSC ability (<250ms). SSC = stretch-shortening cycle.

to assist coaches with providing effective correction of common technical errors and augmentation of performance outcome variables. Finally, effective practical applications of the drop jump are reviewed for suitability within an athletic training program.

TECHNICAL MODEL OF THE DROP JUMP

Drop jumps rely on adequate development of critical biomotor abilities such as maximum strength, rate of force development, SSC function, and leg stiffness (47,89). Consequently, it is necessary to develop high levels of strength in foundational movements, alongside a systematic progression of jump landing-based exercises that gradually and progressively increase the eccentric load placed on the athlete. Flanagan and Comyns (34) have provided suitable activities and progressions within their 4-step progression for developing fast SSC performance. However, even with these foundations in place, technical errors may still occur due to faulty movement patterns that can be acutely corrected by practitioners.

The intensity of a drop jump is determined by the eccentric load (69), which is directly influenced by the duration of the exposure to gravitational acceleration. Therefore, the key variable to manipulate as a means of either reducing or increasing the intensity is drop height. Elevated drop height increases impact velocity, which may subsequently generate greater impact peaks and loading rates if the task exceeds the athlete's eccentric force producing capacities (34,89). To avoid such an eventuality, it is desirable to use an optimal drop height to maximize performance adaptations and minimize the risk of injury. Many studies arbitrarily assign a drop height of 40 cm for all participants (20,39,82), but this might exceed the eccentric strength capabilities of some athletes or fail to reach a threshold for adaptation of others. Several methods have been used to determine the appropriate drop height for training, but few have been experimentally validated. Byrne et al. (17) compared drop jump training with

a height determined by maximum countermovement jump height against the height that achieves optimum reactive strength index (RSI) as suggested by Flanagan and Comyns (34). The optimum RSI method produced a drop height approximately 10 cm lower than the maximum countermovement jump method. Training with a drop height determined by either method produced significant improvements in countermovement jump height and RSI. Therefore, because the RSI method produces a lower drop height but the same training adaptation this might be the best option, as there will be lower impact forces because of the lower drop height. However, the coach's assessment of technical competency should remain as the primary determining factor for the selection of an appropriate drop height. An athlete must demonstrate technical mastery of the exercise at the prescribed drop height; otherwise the intensity should be regressed to avoid unnecessary risk of injury.

To assist practitioners with a conceptual target to direct their athletes toward, a technical model for the drop jump has been provided (Figure 2). The drop jump has been broken down into 5 distinct phases, although there are also some technical factors that should remain consistent throughout the exercise. The hips, knees, and feet should all remain parallel in the frontal plane and therefore, lateral tilt of the pelvis should be minimal. A neutral spine and pelvis position should be maintained throughout the exercise. Finally, a constant fixed gaze should be maintained on a point at head height directly in front of the athlete. It should be noted that there is no arm swing evident in the technical model presented. This is common practice in research studies to try to standardize the height of the center of mass on both landings and to minimize anterior-posterior deviation between the 2 landings. The use of an arm swing has been demonstrated to enhance jump height (33,50). Thus, when working with athletes for performance outcomes, practitioners should allow athletes to practice this skill because it

will be freely available in their sporting context.

EFFECTIVE VERBAL CUEING FOR THE DROP JUMP

Good coaching requires accurate instruction, error identification, relevant and well-timed feedback, with the goal of improving technical proficiency and performance outcomes (19,25,91). There is a growing body of evidence that instructions and feedback used by strength and conditioning coaches have an impact on the acute technical execution and the performance outcomes of a task (1,36,44,68,92,95,96,98). External cues divert the attention to the environment around the athlete and their impact on it, whereas internal cues direct the attention toward the athlete's body, body segments, or body movements (94). An external focus of attention has been shown to augment performance in a number of skills and performance tasks including vertical jumping (93,96). Analogy learning further reduces the amount of explicable information by providing a biomechanical metaphor for a complex motor skill (9,45,51,58). This approach to cueing has been shown to develop performance characteristics that are associated with implicit learning (51). Therefore, strength and conditioning practitioners should attempt to use analogy cues and/or external focus of attention cues when coaching the drop jump, particularly with athletes of a lower training age. Figure 2 provides examples of cues that could be used for the drop jump.

Drop jumping is unique from other jumping tasks in that there are 2 outcome priorities; maximizing jump height and minimizing ground-contact time. Practitioners frequently instruct athletes to jump as high as possible while minimizing the time spent on the ground. This focus on ground-contact time invariably leads to a reduction in jump height, greater peak vertical ground reaction force (VGRF), and greater loading rates because of the stiffer landings usually observed (44,97). In one 6-week






| Phase | Key Points | Common Errors | Corrective Cues |
|---|---|--|--|
|  Step-off | <ol style="list-style-type: none"> 1. The athlete should stand upright on a box with the hands placed on the hips. 2. The movement should be initiated by stepping out from the box with a single leg rather than jumping with both. | Stepping down from or jumping off the box. | <i>"step onto an invisible box"</i> <i>"step out"</i> |
|  Descent | <ol style="list-style-type: none"> 1. As the athlete descends toward the floor, they should prepare for ground contact. 2. Limbs and trunk should be stiffened with the ankle in a neutral position to promote ankle stiffness. 3. A small amount of flexion in the knee and hip should be present. | Excessive forward trunk lean/ looking at the floor. Lack of stiffness in preparation for ground contact. | <i>"Look at a fixed point in front of you"</i> <i>"Be ready to push the floor away immediately"</i> |
|  Contact Phase | <ol style="list-style-type: none"> 1. On ground contact, the feet should be shoulder width apart and the heels of the feet should remain off the floor. 2. The center of mass is likely to fall a small distance during ground contact due to a small amount of hip, knee and ankle flexion and should occur quickly before the movement is rapidly reversed. | Soft landing with excessive knee and hip flexion and very long ground contact times. Poor utilisation of elastic energy and SSC due to lack of preparatory stiffening for impact. Heels collapsing onto floor. Very stiff landing with little hip or knee flexion. Knee valgus | <i>"Bounce like a ball"</i> <i>"Imagine you are on a trampoline or pogo stick"</i> <i>"Try to be quiet on the floor"</i> <i>"Don't squash the grape under your heel"</i> <i>"Bounce like a spring"</i> <i>"Stretch an imaginary band that is around your knees"</i> |
|  Take-off | <ol style="list-style-type: none"> 1. At the point of take-off, the toes should be the final part of the foot to leave the floor. 2. The hip, knee and ankle should all be fully extended as the result of an explosive triple extension in a vertical direction. | Lack of triple extension. Lack of synchronisation of triple extension | <i>"Look over the fence"</i> <i>"Imagine you are being stretched"</i> <i>"Be like a string being pulled tight"</i> |
|  Second Landing | <ol style="list-style-type: none"> 1. Initial contact is made by the forefoot, followed shortly by the heel, meaning weight distribution will move to the rear foot as more of the landing force is absorbed. 2. The athlete should land softly assuming a half-squat position with knees aligned over the toes and feet shoulder distance apart. | Heavy landing with poor force absorption. Poor weight distribution through foot, staying predominantly through the forefoot. There is large horizontal displacement between the first and second landing. | <i>"don't make a sound"</i> <i>"sit onto the chair behind you"</i> <i>"land behind this line"</i> |

Figure 2. Technical model for the drop jump with common errors and example corrective cues.

training study, 3 participants from a training group with a contact time focus were forced to dropout because of tibial pain (97). It was speculated that these dropouts were injured because of low levels of relative leg strength (62). It may be that these findings are more a reflection of inappropriate programming in terms of the prescribed drop height, which was beyond the capabilities of the athlete rather than a direct effect of the cue per se. A focus on contact time when dropping from a drop height that exceeds the eccentric capabilities of the athlete will result in undesirable stiffening strategies such as landing with a much more extended knee and hip and thus placing greater stress on skeletal

structures rather than the muscle-tendon unit.

From a performance perspective, reducing ground-contact time while maintaining jump height is highly desirable as it is a marker of increased power capabilities in the athlete (44). To reduce ground-contact time and maintain jump height, greater VGRF is required in a shorter timeframe to maintain the necessary level of vertical impulse to achieve the same take-off velocity. Consequently, it might be prudent to avoid this attentional focus with athletes who do not have a sufficient level of relative strength (62) because they will be unable to tolerate the loading rates and peak forces effectively, which could elevate their risk of

injury. Previous research has observed significant relationships between the volume of landing sound and the magnitude of the VGRF (87). Because large impact forces place greater stress on soft-tissue structures, it would be efficacious to use cues that encourage the performer to be quiet in both landings involved in the drop jump to avoid undesirably large impact forces during the eccentric portion of ground contact.

Previous literature discussing training progressions to develop SSC function has recommended a progression from tasks that are characterized by a low eccentric demand with a short contact time focus, toward activities involving higher eccentric loads and a greater

focus on increased jump height and reduced ground-contact time (34). These guidelines are intuitive, but it should be noted that a reduced contact time focus might concomitantly elevate ground reaction forces on impact; therefore, the task provided to the athlete must involve a level of eccentric load that can be tolerated to optimize training adaptations (44). This serves as another important reminder of conservative progression in the use of plyometric training. Regardless of whether the focus is directed toward jump height or ground-contact time, the literature is congruent in its support for maintaining an external focus of attention.

Finally, it should be noted that the effectiveness of all cues will vary between individuals. What works for 1 athlete may not work for another. It is also important to be aware of the potential negative effects of cues if they are taken too literally or to extremes by the athlete. For example, instructing an athlete to “land quietly” during the initial landing of a drop jump is intended to reduce excessive impact peaks and improve SSC utilization; however, if taken too literally by the athlete, they may use excessive magnitudes of hip and knee flexion and let the heels contact the floor to give themselves more time to absorb force. As a coach, previous awareness of these undesirable outcomes is necessary to allow early detection and implementation of a different cueing strategy that is more easily and accurately interpreted by the athlete. Because of the high degree of individual variability in cueing effectiveness, it is essential that a coach establishes meaning with their athletes and has a battery of different cues for prompting the same movement when working with a wide range of individuals.

WHY USE THE DROP JUMP?

PERFORMANCE

When selecting appropriate exercises, their ability to transfer adaptations into improved execution of a sporting task

should be considered. The magnitude, direction and rate of force production, the nature of muscular contractions, and the energy systems used are all important factors in determining the degree of training transfer between an exercise and a sports movement (78). Most human movements use the SSC rather than relying purely on eccentric, concentric, or isometric actions (46). This is particularly true of locomotive tasks such as walking, running, skipping, hopping, and jumping (63,74). These actions typically involve limited time for force production because of brief ground-contact times and some element of rebounding, with the main propulsive action being simultaneous triple extension of the ankle, knee, and hip joints. Similarly, drop jumps also involve SSC muscle actions with particular emphasis on eccentric overload during the yielding phase where rapid triple flexion takes place, and this is followed by triple extension to propel the body into the air. This sequence is replicable of the time frames afforded in human locomotive tasks such as jumping and ground-contact times during sprinting and change-of-direction tasks (14,57,61).

In practice, drop jumps are rarely used in isolation but rather as a component of an athlete's training program. However, the performance enhancing effects of plyometrics are well documented (6,55,72,80) and drop jumps in isolation have been shown to improve countermovement, squat and drop jump performance (39,59,97), 5, 20, and 40 m linear sprinting ability (20,73), 505 and *t* test change-of-direction ability (82,85), and running economy (5,10). Drop jumping lies at the most intense end of the plyometric training spectrum (34,43,52), and with correct implementation is therefore a suitable exercise in the training programs of well-trained athletes seeking fast SSC development.

POSTACTIVATION POTENTIATION

In addition to providing a chronic training stimulus, drop jumps have also

been used in an attempt to acutely improve athletic performance in a variety of tasks. Postactivation potentiation (PAP) is an acute facilitation of enhanced muscle performance through the use of a preceding activity before the execution of another task (18,84). The inclusion of low volume (2–6 repetitions) drop jumps at the end of a warm-up protocol has been shown to improve subsequent sprint performance (15,18), vertical jump ability (15), and 1 repetition maximum back squat (16). Conversely, other studies have found no beneficial acute vertical jump performance effects after drop jumps used as a potentiating stimulus (31,81). This may be due to strength levels, fiber type distribution, and training age, all of which have been postulated as confounding factors (83) impairing the effectiveness of the SSC (47). To this end, a certain level of physical competency in utilization of the fast SSC may be necessary to elicit a PAP response from drop jumps. Furthermore, careful consideration needs to be given to the selection of drop height on an individual basis to allow eccentric loading within athlete's tolerance levels and to maximize the probability of a PAP response.

TRAINING MONITORING TOOL

Because of the reliance on both neurological and muscular mechanisms during drop jumps, the RSI has been used to assess the levels of neuromuscular fatigue and quantify readiness to train (12,49,56,67). RSI is the ratio of jump height to ground-contact time (34) and has shown to be reliable in numerous studies (32,35,60). A good RSI score is achieved through maximizing jump height, while minimizing ground-contact time. Executing this effectively requires good SSC function and high levels of leg and ankle stiffness (2).

Drop jump performance is susceptible to fatigue after marathon running (3) and simulated soccer activity (65). In both instances, greater reductions in peak VGRF between the impact peak and the propulsive peak were evident,

which suggests a decrement in SSC function because of fatigue. Links between fatigue and injury risk are well documented in team sports (41,65,66), with most injuries occurring toward the end of playing time (30,88) and during periods of the season where fixtures are congested and players experience the most cumulative fatigue (8,37,70). RSI has been shown to be sensitive to increased workloads and fatigue in elite rugby union players and youth soccer players during tournament match play (4,41). Therefore, this metric could be considered a useful assessment tool to monitor neuromuscular fatigue with athletes, provided they are familiar with drop jump training and technique.


Based on the cumulative body of evidence, it seems that the drop jump has many uses for training and monitoring athletes. Drop jumps can be used as an effective fast SSC development exercise in well-trained athletes who possess sufficient levels of strength and are able to tolerate high eccentric forces. The drop jump also offers promise as an injury-risk screening tool, with kinematic variables showing good predictive ability; however, further research is warranted to explore the link between injury risk and a range of kinetic variables that assess relevant forces and rates of loading related to the mechanisms of traumatic injury. Finally, the drop jump can be used as an objective daily readiness-to-train measurement tool to optimize training loads, reduce the risk of nonfunctional overreaching, and injury due to athletes training in a fatigued state.

SUMMARY

The drop jump is a cornerstone of plyometric athletic training programs but has other diverse applications including injury-risk screening and monitoring of training and neuromuscular readiness. The drop jump requires well-developed function of the SSC and provides a unique challenge to the athlete in comparison with other SSC activities such as

countermovement jumps because of the greater eccentric load, elevated power output, and magnitude of impact forces. Proficient execution of the drop jump requires high levels of strength, in addition to effective and safe movement control. The latter can be developed through effective coaching with an emphasis on external cueing as part of a periodized training program, and with progression determined by technical competency. This article has also proposed a technical model for the drop jump and guidelines to correct common technical errors through effective cueing, which has the potential to improve performance and reduce the likelihood of musculoskeletal injury.

Conflicts of Interest and Source of Funding: The authors report no conflicts of interest and no source of funding.




University.

Jason S. Pedley is a lecturer in strength and conditioning and coach within the Youth Physical Development Centre, Cardiff Metropolitan



Centre, Cardiff Metropolitan University.

Rhodri S. Lloyd is a senior lecturer in strength and conditioning and chair of the Youth Physical Development



Hospital.

Paul Read is a strength and conditioning coach and clinical research scientist, Aspetar Orthopaedic and Sports Medicine



University.

Isabel S. Moore is a lecturer in sport and exercise medicine and programme director of the masters in sport and exercise medicine course, Cardiff Metropolitan



University.

Jon L. Oliver is a reader in Applied Paediatric Exercise Science and co-founder of the Youth Physical Development Group, Cardiff Metropolitan

REFERENCES

1. Abdollahipour R, Wulf G, Psotta R, and Nieto MP. Performance of gymnastics skill benefits from an external focus of attention. *J Sports Sci* 33: 1807–1813, 2015.
2. Arampatzis A, Schade F, Walsh M, and Brüggemann GP. Influence of leg stiffness and its effect on myodynamic jumping performance. *J Electromyogr Kinesiol* 11: 355–364, 2001.
3. Avela J, Kyröläinen H, Komi PV, and Rama D. Reduced reflex sensitivity persists several days after long-lasting stretch-shortening cycle exercise. *J Appl Physiol* 86: 1292–1300, 1999.
4. Beattie K and Flanagan EP. Establishing the reliability and meaningful change of the drop-jump reactive strength index. *J Aust Strength Cond* 23: 12–18, 2015.
5. Beattie K, Kenny IC, Lyons M, and Carson BP. The effect of strength training on performance in endurance athletes. *Sport Med* 44: 845–865, 2014.
6. Bedoya AA, Miltenberger MR, and Lopez RM. Plyometric training effects on athletic performance in youth soccer athletes: A systematic review. *J Strength Cond Res* 29: 2351–2360, 2015.
7. Behrens M, Mau-Moeller A, Mueller K, Heise S, Gube M, Beuster N, Herlyn PK, Fischer DC, and Bruhn S. Plyometric

- training improves voluntary activation and strength during isometric, concentric and eccentric contractions. *J Sci Med Sport* 19: 1–7, 2015.
8. Bengtsson H, Ekstrand J, and Häggglund M. Muscle injury rates in professional football increase with fixture congestion: An 11-year follow-up of the UEFA champions league injury study. *Br J Sports Med* 47: 743–747, 2013.
9. Bennett SJ. Implicit learning: Should it be used in practice? *Int J Sport Psychol* 31: 542–646, 2000.
10. Berryman N, Maurel D, and Bosquet L. Effect of plyometric vs. dynamic weight training on the energy cost of running. *J Strength Cond Res* 24: 1818–1825, 2010.
11. Blagrove R. *Strength and Conditioning for Endurance Running*. Marlborough: Wiltshire: Crowood Press, 2015.
12. Blazeovich A. The stretch-shortening cycle (SSC). In: *Strength and Conditioning: Biological Principles and Practical Applications*. Cardinale M, Newton RU, and Nosaka K, eds. Oxford, United Kingdom: Wiley-Blackwell, 2011. pp. 209–222.
13. Bobbert MF, Huijij PA, and van Ingen Schenau GJ. Drop jumping I: The influence of jumping technique on the biomechanics of jumping. *Med Sci Sport Exerc* 19: 332–338, 1987.
14. Bobbert MF, Huijij PA, and van Ingen Schenau GJ. Drop jumping II: The influence of dropping height on the biomechanics of drop jumping. *Med Sci Sport Exerc* 19: 339–346, 1987.
15. Bomfim Lima JC, Marin DP, Barquilha G, Da Silva LO, Puggina EF, Pithon-Curi TC, and Hirabara S. Acute effects of drop jump potentiation protocol on sprint and countermovement vertical jump performance. *Hum Mov* 12: 324–330, 2011.
16. Bullock N and Comfort P. An investigation into the acute effects of depth jumps on maximal strength performance. *J Strength Cond Res* 25: 3137–3141, 2011.
17. Byrne P, Moran K, Rankin P, and Kinsella S. A comparison of methods used to identify “optimal” drop height for early phase adaptations in depth jump training. *J Strength Cond Res* 24: 2050–2055, 2010.
18. Byrne P, Kenny J, and O’ Rourke B. Acute potentiating effect of depth jumps on sprint performance. *J Strength Cond Res* 28: 610–615, 2014.
19. Chambers F. Learning theory for effective learning in practice. In: *Sport Pedagogy: An Introduction for Teaching and Coaching*. Armour K, ed. Harlow, United Kingdom: Pearson, 2011. pp. 39–52.
20. Chelly MS, Ghenem MA, Abid K, Hermassi S, Tabka Z, and Shephard RJ. Effects of in-season short-term plyometric training program on leg power, jump-and sprint performance of soccer players. *J Strength Cond Res* 24: 2670–2676, 2010.
21. Chu DA. *Jumping into Plyometrics*. Champaign, IL: Human Kinetics, 1992.
22. Chu DA. Explosive power. In: *High-performance Sports Conditioning*. Foran B, ed. Champaign, IL: Human Kinetics, 2001. pp. 83–98.
23. Cormie P, McBride JM, and McCaulley GO. Power-time, force-time, and velocity-time curve analysis of the countermovement jump: Impact of training. *J Strength Cond Res* 23: 177–186, 2009.
24. Cowling EJ, Steele JR, and McNair PJ. Effect of verbal instructions on muscle activity and risk of injury to the anterior cruciate ligament during landing. *Br J Sports Med* 37: 126–130, 2003.
25. Cross N. Coaching effectiveness. In: *Coaching Process: Principles and Practice for Sport*. Cross N and Lyle J, eds. Oxford, United Kingdom: Butterworth-Heinemann, 1999.
26. Duehring MD, Feldmann CR, and Ebben WP. Strength and conditioning practices of United States high school strength and conditioning coaches. *J Strength Cond Res* 23: 2188–2203, 2009.
27. Ebben W and Blackard P. Strength and conditioning practices of National Football League teams. *J Strength Cond Res* 15: 48–58, 2001.
28. Ebben WP, Carroll RM, and Simenz CJ. Strength and conditioning practices of National Hockey League strength and conditioning coaches. *J Strength Cond Res* 18: 889–897, 2004.
29. Ebben WP, Hintz MJ, and Simenz CJ. Strength and conditioning practices of Major League Baseball strength and conditioning coaches. *J Strength Cond Res* 19: 538–546, 2005.
30. Ekstrand J, Häggglund M, and Walden M. Injury incidence and injury patterns in professional football: The UEFA injury study. *Br J Sports Med* 45: 553–558, 2011.
31. Faulkinbury KJ, Stieg JL, Tran TT, Brown LE, Coburn JW, and Judelson DA. Effects of depth jump vs. box jump warm-ups on vertical jump in collegiate vs. club female volleyball players. *Med Sport* 15: 103–106, 2011.
32. Feldman C, Weiss L, Ferreira L, Schilling BK, and Hammond K. Reactive strength index and ground contact time: Reliability, precision, and association with drop vertical jump displacement. *J Strength Cond Res* 25: S1, 2011.
33. Feltner ME, Fraschetti DJ, and Crisp RJ. Upper extremity augmentation of lower extremity kinetics during countermovement vertical jumps. *J Sports Sci* 17: 449–466, 1999.
34. Flanagan EP and Comyns TM. The use of contact time and the reactive strength index to optimize fast stretch-shortening cycle training. *Strength Cond J* 30: 32–38, 2008.
35. Flanagan EP, Ebben WP, and Jensen RL. Reliability of the reactive strength index and time to stabilization during depth jumps. *J Strength Cond Res* 22: 1677–1682, 2008.
36. Freudenheim AM, Wulf G, Madureira F, Pasetto SC, and Corrêa UC. An external focus of attention results in greater swimming speed. *Int J Sport Sci Coach* 5: 533–542, 2010.
37. Gabbett TJ. Influence of injuries on team playing performance in rugby league. *J Sci Med Sport* 7: 340–346, 2004.
38. Gee TI, Olsen PD, Berger NJ, Golby J, and Thompson KG. Strength and conditioning practices in rowing. *J Strength Cond Res* 25: 668–682, 2011.
39. Gehri DJ, Ricard MD, Kleiner DM, and Kirkendall DT. A comparison of plyometric training techniques for improving vertical jump ability and energy production. *J Strength Cond Res* 12: 85, 1998.
40. Goodwin JE and Jeffreys I. Plyometric Training: Theory and practice. In: *Strength and Conditioning for Sports Performance*. Jeffreys I and Moody J, eds. London, United Kingdom: Routledge, 2016. pp. 304–340.
41. Hamilton D. Drop jumps as an indicator of neuromuscular fatigue and recovery in elite youth soccer athletes following tournament match play. *J Aust Strength Cond* 17: 3–8, 2009.
42. Hartmann J and Tunnemann H. *Fitness and Strength Training for All Sports*. Toronto, Ontario: Sport Books Publisher, 2001.
43. Jarvis MM, Graham-Smith P, and Comfort P. A methodological approach to quantifying plyometric intensity. *J Strength Cond* 30: 2522–2532, 2016.
44. Khuu S, Musalem LL, and Beach TA. Verbal instructions acutely affect drop vertical jump biomechanics- implications for athletic performance and injury risk assessments. *J Strength Cond Res* 29: 2816–2826, 2015.

45. Komar J, Chow JY, Chollet D, and Seifert L. Effect of analogy instructions with an internal focus on learning. *J Appl Sport Psychol* 26: 17–32, 2014.
46. Komi PV. Stretch-shortening cycle: A powerful model to study normal and fatigued muscle. *J Biomech* 33: 1197–1206, 2000.
47. Komi PV. Stretch-shortening cycle. In: *Strength and Power in Sport: Volume III of the Encyclopaedia of Sports Medicine- an IOC Medical Commission Publication*. Komi PV, ed. Oxford, United Kingdom: Blackwell Science, 2003. pp. 184–202.
48. Komi PV and Bosco C. Utilization of stored elastic energy in leg extensor muscles by men and women. *Med Sci Sports* 10: 261–265, 1978.
49. Kuitunen S, Avela J, Kyröläinen H, Nicol C, and Komi PV. Acute and prolonged reduction in joint stiffness in humans after exhausting stretch-shortening cycle exercise. *Eur J Appl Physiol* 88: 107–116, 2002.
50. Lees A, Vanrenterghem J, and Clercq DD. Understanding how an arm swing enhances performance in the vertical jump. *J Biomech* 37: 1929–1940, 2004.
51. Liao C and Masters RSW. Analogy learning : A means to implicit motor learning. *J Sports Sci* 19: 307–319, 2001.
52. Lloyd RS, Meyers RW, and Oliver JL. The natural development and trainability of plyometric ability during childhood. *Strength Cond J* 33: 23–32, 2011.
53. Makaruk H, Porter JM, Czaplicki A, Sadowski J, and Sacewicz T. The role of attentional focus in plyometric training. *J Sports Med Phys Fitness* 52: 319–327, 2012.
54. Markovic G and Mikulic P. Neuro-musculoskeletal and performance adaptations to lower-extremity plyometric training. *Sport Med* 40: 859–895, 2010.
55. Markovic G and Newton RU. Does plyometric training improve vertical jump height? A meta-analytical review-commentary. *Br J Sports Med* 41: 349–355, 2007.
56. Markwick WJ, Bird SP, Tufano JJ, Seitz LB, and Haff GG. The intraday reliability of the reactive strength index calculated from a drop jump in professional men's basketball. *Int J Sports Physiol Perform* 10: 482–488, 2015.
57. Marshall BM and Moran KA. Which drop jump technique is most effective at enhancing countermovement jump ability, "countermovement" drop jump or "bounce" drop jump? *J Sports Sci* 31: 1368–1374, 2013.
58. Masters RSW. Theoretical aspects of implicit learning in sport. *Int J Sport Psychol* 31: 530–541, 2000.
59. McClenton LS, Brown LE, Coburn JW, and Kersey RD. The effect of short-term VertiMax vs. depth jump training on vertical jump performance. *J Strength Cond Res* 22: 321–325, 2008.
60. McClymont D. Use of the reactive strength index (RSI) as an indicator of plyometric training conditions. In: *Science and Football V: Proceedings of the 5th World Congress on Science and Football*. Reilly T, Cabri J, and Araujo D, eds. London: Routledge, 2005. pp. 423–430.
61. Morin JB, Slawinski J, Dorel S, De Villareal ES, Couturier A, Samozino P, Brughelli M, and Rabita G. Acceleration capability in elite sprinters and ground impulse: Push more, brake less?. *J Biomech* 48: 1–6, 2015.
62. Newton RU, Young WB, Kraemer WJ, and Byrne C. Effects of drop jump height and technique on ground reaction force with possible implication for injury. *Sport Med Train Rehabil* 10: 83–93, 2001.
63. Nigg BM, MacIntosh BR, and Mester J. *Biomechanics and Biology of Movement*. Champaign, IL: Human Kinetics, 2000.
64. Noyes FR, Barber-Westin SD, Fleckenstein C, Walsh C, and West J. The drop-jump screening test: Difference in lower limb control by gender and effect of neuromuscular training in female athletes. *Am J Sports Med* 33: 197–207, 2005.
65. Oliver J, Armstrong N, and Williams C. Changes in jump performance and muscle activity following soccer-specific exercise. *J Sports Sci* 26: 141–148, 2008.
66. Oliver J, De Ste Croix M, Cone J, and Greig M. Altered neuromuscular control of leg stiffness following soccer-specific exercise. *Eur J Appl Physiol* 114: 2241–2249, 2014.
67. Padua DA, Arnold BL, Perrin DH, Gansneder BM, Carcia CR, and Granata KP. Fatigue, vertical leg stiffness, and stiffness control strategies in males and females. *J Athl Train* 41: 294–304, 2006.
68. Porter J, Strowski E, and Nolan R. Standing long-jump performance is enhanced when using an external focus of attention. *J Strength Cond Res* 24: 1746–1750, 2010.
69. Potach DH and Chu DA. Plyometric training. In: *Essentials of Strength Training and Conditioning*. Baechle TR and Earle RW, eds. Champaign, IL: Human Kinetics, 2008. pp. 413–456.
70. Price RJ, Hawkins RD, Hulse MA, and Hodson A. The Football Association medical research programme: An audit of injuries in academy youth football. *Br J Sports Med* 38: 466–471, 2004.
71. Rathleff MS, Richter C, Brushej C, Bencke J, Bandholm T, Hölmich P, and Thorborg K. Increased medial foot loading during drop jump in subjects with patellofemoral pain. *Knee Surgery, Sport Traumatol Arthrosc* 22: 2301–2307, 2014.
72. Sáez-Sáez de Villarreal E, Requena B, and Newton RU. Does plyometric training improve strength performance? A meta-analysis. *J Sci Med Sport* 13: 513–522, 2010.
73. Saez de Villarreal ES, González-Badillo JJ, and Izquierdo M. Low and moderate plyometric training frequency produces greater jumping and sprinting gains compared with high frequency. *J Strength Cond Res* 22: 715–725, 2008.
74. Sawicki GS, Lewis CL, and Ferris DP. It pays to have a spring in your step. *Exerc Sport Sci Rev* 37: 130–138, 2009.
75. Schmidtleicher D. Training for power events. In: *The Encyclopedia of Sports Medicine*. Komi PV, ed. Oxford, United Kingdom: Blackwell Science, 1992. pp. 381–395.
76. Schmitz RJ, Cone JC, Tritsch AJ, Pye ML, Montgomery MM, Henson RA, and Shultz SJ. Changes in drop-jump landing biomechanics during prolonged intermittent exercise. *Sports Health* 6: 128–135, 2014.
77. Sheppard J. Optimising training for jumping and landing. In: *High-performance Training for Sports*. Joyce D and Lewindon D, eds. Champaign, IL: Human Kinetics, 2014. pp. 167–183.
78. Siff MC. *Supertraining*. First. Denver, CO: Elite Fitness Systems, 2000.
79. Simenz CJ, Dugan CA, and Ebben WP. Strength and conditioning practices of National Basketball Association strength and conditioning coaches. *J Strength Cond Res* 19: 495–504, 2005.
80. Spurr RW, Murphy AJ, and Watsford ML. The effect of plyometric training on distance running performance. *Eur J Appl Physiol* 89: 1–7, 2003.
81. Stieg JL, Faulkinbury KJ, Tran TT, Brown LE, Coburn JW, and Judelson DA. Acute effects of depth jump volume on vertical jump performance in collegiate women soccer players. *Kinesiology* 43: 25–30, 2011.
82. Thomas K, French D, and Hayes PR. The effect of two plyometric training techniques

- on muscular power and agility in youth soccer players. *J Strength Cond Res* 23: 332–335, 2009.
83. Tillin NA and Bishop D. Factors modulating post-activation potentiation and its effect on performance of subsequent explosive activities. *Sport Med* 39: 147–166, 2009.
 84. Turner AN, Bellhouse S, Kilduff LP, and Russell M. Postactivation potentiation of sprint acceleration performance using plyometric exercise. *Strength Cond J* 29: 343–350, 2015.
 85. Vácsi M, Tollár J, Meszler B, Juhász I, and Karsai I. Short-term high intensity plyometric training program improves strength, power and agility in male soccer players. *J Hum Kinet* 36: 17–26, 2013.
 86. Verkhoshansky Y and Verkhoshansky N. *Special Strength Training: Manual for Coaches*. Rome, IT: Verkhoshansky SSTM, 2011.
 87. Wernli K, Ng L, Phan X, Davey P, and Grisbrook T. The relationship between landing sound, vertical ground reaction force and kinematics of the lower limb during drop landings in healthy males. *J Orthop Sport Phys Ther* 46: 1–22, 2016.
 88. Williams S, Trewartha G, Kemp S, and Stokes K. A meta-analysis of injuries in senior men's professional Rugby Union. *Sport Med* 43: 1043–1055, 2013.
 89. Wilson JM and Flanagan EP. The role of elastic energy in activities with high force and power requirements: A brief review. *J Strength Cond Res* 22: 1705–1715, 2008.
 90. Winwood PW, Keogh JW, and Harris NK. The strength and conditioning practices of strongman competitors. *J Strength Cond Res* 25: 3118–3128, 2011.
 91. Wulf G. *Attention and Motor Skill Learning*. Champaign, IL: Human Kinetics, 2007.
 92. Wulf G. Attentional focus effects in balance acrobats. *Res Q Exerc Sport* 79: 319–325, 2008.
 93. Wulf G, Dufek JS, Lozano L, and Pettigrew C. Increased jump height and reduced EMG activity with an external focus. *Hum Mov Sci* 29: 440–448, 2010.
 94. Wulf G and Prinz W. Directing attention to movement effects enhances learning: A review. *Psychon Bull Rev* 8: 648–660, 2001.
 95. Wulf G and Su J. An external focus of attention enhances golf shot accuracy in beginners and experts. *Res Q Exerc Sport* 78: 384–389, 2007.
 96. Wulf G, Zachry T, Granados C, and Dufek JS. Increases in jump-and-reach height through an external focus of attention. *Int J Sport Sci Coach* 2: 275–284, 2007.
 97. Young W, Wilson C, and Byrne C. A comparison of drop jump training methods: Effects on leg extensor strength qualities and jumping performance. *Int J Sports Med* 20: 295–303, 1999.
 98. Young WB, Pryor JF, and Wilson GJ. Effect of instructions on characteristics of countermovement and drop jump performance. *J Strength Cond Res* 9: 232–236, 1995.



APPLY YOUR

STRENGTH

Devote your time to helping others by assessing, motivating, educating, and training individuals for the primary goal of improving their personal health and fitness goals! Differentiate yourself from the competition and begin your path to becoming an NSCA-Certified Personal Trainer® (NSCA-CPT®) today.

Learn more about the NSCA-CPT certification at
NSCA.com/getCPT

